

<b>Citation</b>	T. Odenthal, T. Heck, B. Smeets, P. Van Liedekerke, H. Ramon, H. Van Oosterwyck <b>Coupling of a cell migration model with an (N)SPH substrate to investigate dynamic generation of cellular tractions</b>
<b>Archived version</b>	Author manuscript: the content is identical to the content of the published paper, but without the final typesetting by the publisher
<b>Conference</b>	4th International Conference on Particle-based Methods, 28-30 September 2015, Barcelona, Spain
<b>Author contact</b>	Hans.vanoosterwyck@kuleuven.be
<b>Acknowledgements</b>	The research leading to these results has received funding from the European Research Council under the European Union's Seventh Framework Programme (FP7/2007-2013)/ ERC Grant Agreement n° 308223), FWO-Vlaanderen (doctoral fellowship of TH; G.0821.13) and the Agency for Innovation by Science and Technology in Flanders (IWT) (doctoral fellowship of BS) .



# Coupling of a cell migration model with an (N)SPH substrate to investigate dynamic generation of cellular tractions

Tim Odenthal\*, Tommy Heck†, Bart Smeets\*, Paul Van Liedekerke#, Herman Ramon\*, Hans Van Oosterwyck†

\* MeBioS, KU Leuven, Kasteelpark Arenberg 30,  
3001 Heverlee, Belgium  
e-mail: Tim.Odenthal@kuleuven.be, web page: <http://www.mebios.be>

† Biomechanics Section, KU Leuven, Celestijnenlaan 300,  
3001 Heverlee, Belgium

# INRIA (Paris) - Domaine de Voluceau-Rocquencourt, B.P. 105  
8153 Le Chesnay Cedex, France

## ABSTRACT




Figure 1: Cell migrating to the right on a visco-elastic NSPH substrate. Von Mises stress is shown in substrate colour; highest stresses appear under the lamellipodium in the front.

Cell migration is crucial for tissue development, maintenance and repair. To migrate, cells apply traction on the extra-cellular matrix (ECM) and degrade it. At the same time, the response of the ECM is an important cue for the cell to modulate mechanotransduction processes.

To investigate cell-ECM interaction in the context of active spreading and migration, a previously developed deformable cell model [1] has been extended by (amongst others) including a protrusive pressure at the leading edge of the cell. In this model, the membrane/actin cortex of the cell is discretized into deformable rounded triangles, over which contact pressures including adhesion are integrated between the cell and the ECM. The matrix represents a typical substitute ECM in *in vitro* experiments that is continuous at the cellular level and has tunable physical and mechanical properties. To be able to model the possibly large deformations and degradation of this hydrogel, we chose the Non-inertial, Smoothed-Particle Hydrodynamics (NSPH) [2] method to avoid frequent re-meshing that is typically required in finite element methods.

We will discuss the extent to which this method can capture typical hydrogel properties by comparing with FEM simulations of a simplified system. In a second step, we will highlight the effects of hydrogel properties as well as the properties of the hydrogel-cell interaction on active spreading and migration of the deformable cell. In particular, the model is able to capture stress magnitudes and distributions which correspond to typical traction force microscopy results for migratory epithelial cells, see Figure 1.

## REFERENCES

- [1] Odenthal, T., Smeets, B., et al. (2013). Analysis of initial cell spreading using mechanistic contact formulations for a deformable cell model. *PLoS Computational Biology*, 9 (10), e1003267.
- [2] Van Liedekerke, P., et al. (2013). Solving microscopic flow problems using Stokes equations in SPH. *Comp. Phys. Comm.*, 184 (7), 1686-1696.